

Linear Codes from the Axiomatic Viewpoint

Jay A. Wood

Department of Mathematics
Western Michigan University
<http://homepages.wmich.edu/~jwood/>

Noncommutative rings and their applications, IV
University of Artois, Lens
June 9, 2015

4. MacWilliams extension theorem for other weights

- ▶ Homogeneous and egalitarian weights
- ▶ Symmetrized weight compositions
- ▶ General weight: reducing to symmetrized weight compositions
- ▶ Weights with maximal symmetry
- ▶ Lee and Euclidean weights on $\mathbb{Z}/N\mathbb{Z}$

Notation

- ▶ Let R be a finite associative ring with 1.
- ▶ Let A be a finite unital left R -module: the **alphabet**.
- ▶ Let $w : A \rightarrow \mathbb{Q}$ be a **weight**: $w(0) = 0$. Extend to A^n by

$$w(a_1, \dots, a_n) = \sum_{i=1}^n w(a_i).$$

Symmetry groups

- ▶ Recall the **symmetry groups** of w :

$$G_{\text{lt}} = \{u \in \mathcal{U}(R) : w(ua) = w(a), a \in A\},$$

$$G_{\text{rt}} = \{\phi \in GL_R(A) : w(a\phi) = w(a), a \in A\}.$$

- ▶ $\mathcal{U}(R)$ is the group of units of R , and $GL_R(A)$ is the group of invertible R -linear homomorphisms $A \rightarrow A$.
- ▶ Recall that I will usually write homomorphisms of left modules on the right side.

Orbit spaces

- ▶ For an information module M , recall the **orbit spaces**:

$$\mathcal{O} = G_{\text{lt}} \backslash M$$

$$\mathcal{O}^{\#} = \text{Hom}_R(M, A) / G_{\text{rt}}$$

W -map

- ▶ The W -**map** is

$$W : F_0(\mathcal{O}^\#, \mathbb{Q}) \rightarrow F_0(\mathcal{O}, \mathbb{Q}).$$

- ▶ For $x \in M$,

$$W(\eta)(x) = \sum_{[\lambda] \in \mathcal{O}^\#} w(x\lambda)\eta([\lambda]).$$

Using generating character to define a weight

- ▶ Suppose the alphabet A admits a generating character ρ : $\text{Soc}(A)$ cyclic.
- ▶ Fix a subgroup $U \subseteq GL_R(A)$.
- ▶ Define a weight $w_U : A \rightarrow \mathbb{C}$:

$$w_U(a) = 1 - \frac{1}{|U|} \sum_{\phi \in U} \rho(a\phi), \quad a \in A.$$

Properties of w_U

- ▶ $w_U(0) = 0$.
- ▶ Re-index: $U \subseteq G_{\text{rt}}(w_U)$.
- ▶ By the summation formulas and ρ generating: for any nonzero left R -submodule $B \subseteq A$, and any $a_0 \in A$,

$$\sum_{b \in B} w_U(a_0 + b) = |B|.$$

- ▶ We say that w_U is **egalitarian** on cosets of B .
- ▶ $w_{GL_R(A)}$ is called **homogeneous**: Constantinescu, Heise, Greferath, Schmidt, Honold, Nechaev.

w_U has EP

- ▶ Suppose $w_U(x\Lambda) = w_U(xN)$ for all $x \in M$.
- ▶ Equation of characters: for all $x \in M$,

$$\sum_{i=1}^n \sum_{\phi \in U} \rho(x\lambda_i\phi) = \sum_{j=1}^n \sum_{\psi \in U} \rho(x\nu_j\psi).$$

- ▶ Use linear independence of characters: for $j = 1$, $\psi = \text{id}_A$, there exist $i = \sigma(1)$ and $\phi_1 \in U$ with $\rho(x\lambda_{\sigma(1)}\phi_1) = \rho(x\nu_1)$ for all $x \in M$.
- ▶ ρ generating: $\nu_1 = \lambda_{\sigma(1)}\phi_1$. Inner sums agree, reduce outer sum, and continue by induction.

Symmetrized weight composition

- ▶ This time, no weight. Just ring R , alphabet A , and a subgroup $G \subseteq GL_R(A)$.
- ▶ Define an equivalence relation from the right action of G on A : for $a, b \in A$, $a \sim b$ if $b = a\phi$ for some $\phi \in G$. Denote equivalence class of a by $[a]$.
- ▶ For $a \in A$ and $x = (x_1, \dots, x_n) \in A^n$, define the **symmetrized weight composition** (swc) by

$$\text{swc}_{[a]}(x) = |\{i : x_i \in [a]\}|.$$

- ▶ Example: $R = A = \mathbb{Z}/4\mathbb{Z}$, $G = \{\pm 1\}$.

swc has EP

- ▶ Assume A has generating character ρ (cyclic socle).
- ▶ Suppose $C_1, C_2 \subseteq A^n$ are two left R -linear codes.
- ▶ Suppose $f : C_1 \rightarrow C_2$ is a linear isomorphism of R -modules that preserves swc:

$$\text{swc}_{[a]}(xf) = \text{swc}_{[a]}(x), \quad a \in A, x \in C_1.$$

- ▶ Then f extends to a G -monomial transformation.

Proof (a)

- ▶ Result dates from 1997, but we will use the local-global idea of Barra, Gluesing-Luerssen (2014). This is joint work with N. El Garem and N. Megahed.
- ▶ As before, view preservation of swc in terms of $\Lambda, N : M \rightarrow A^n$:

$$\text{swc}_{[a]}(x\Lambda) = \text{swc}_{[a]}(xN), \quad a \in A, x \in M.$$

- ▶ Local: for each $x \in M$, there exist a permutation σ_x and elements $\phi_{1,x}, \dots, \phi_{n,x} \in G$ with $x\nu_i = x\lambda_{\sigma_x(i)}\phi_{i,x}$.

Proof (b)

- ▶ Local to global: apply $\phi \in G$ and ρ , then sum over ϕ and i . For every $x \in M$:

$$\sum_{i=1}^n \sum_{\phi \in G} \rho(x\nu_i\phi) = \sum_{i=1}^n \sum_{\phi \in G} \rho(x\lambda_{\sigma_x(i)}\phi_{i,x}\phi).$$

- ▶ Dependence on x disappears! For all $x \in M$:

$$\sum_{i=1}^n \sum_{\phi \in G} \rho(x\nu_i\phi) = \sum_{i=1}^n \sum_{\phi \in G} \rho(x\lambda_i\phi).$$

- ▶ Proceed as before to get G -monomial transformation.

General weight: reducing to swc

- ▶ Now include a weight w . Suppose alphabet A has cyclic socle. Form swc using $G = G_{\text{rt}}(w)$.
- ▶ For any $b = (b_1, \dots, b_n) \in A^n$,

$$w(b) = \sum_{i=1}^n w(b_i) = \sum_{[a] \in A/G_{\text{rt}}} w(a) \text{swc}_{[a]}(b).$$

- ▶ For a scalar multiple $rb \in A^n$, $r \in R$:

$$w(rb) = \sum_{[a] \in A/G_{\text{rt}}} w(ra) \text{swc}_{[a]}(b).$$

- ▶ $w(rb)$ depends only on class $[r] \in G_{\text{lt}} \setminus R$.

Sufficient condition for EP for w

- ▶ Form matrix \mathcal{A} with rows indexed by nonzero $[r] \in G_{\text{lt}} \setminus R$ and columns indexed by nonzero $[a] \in A/G_{\text{rt}}$:

$$\mathcal{A}_{[r],[a]} = w(ra).$$

- ▶ If matrix \mathcal{A} has trivial right nullspace, then alphabet A has EP for w .
- ▶ When $A = R$ is commutative, \mathcal{A} is square. Condition is $\det \mathcal{A} \neq 0$.

Proof (a)

- ▶ Suppose $f : C_1 \rightarrow C_2$ is an isomorphism of R -modules and that f is a linear isometry with respect to w . Codes are given by $\Lambda : M \rightarrow A^n$ and $N : M \rightarrow A^n$, as usual.
- ▶ Isometry: $w(x\Lambda) = w(xN)$, for all $x \in M$.
- ▶ For every $x \in M$, $r \in R$:

$$\begin{aligned}
 0 &= w(rx\Lambda) - w(rxN) \\
 &= \sum_{[a] \in A/G_{\text{rt}}} w(ra) \{ \text{swc}_{[a]}(x\Lambda) - \text{swc}_{[a]}(xN) \}
 \end{aligned}$$

Proof (b)

- ▶ The condition on matrix \mathcal{A} implies $\text{swc}_{[a]}(x\Lambda) = \text{swc}_{[a]}(xN)$ for every $a \in A$ and $x \in M$.
- ▶ This means that $f : C_1 \rightarrow C_2$ preserves swc.
- ▶ Apply EP for swc to conclude that f extends to a G_{rt} -monomial transformation.
- ▶ This result has been used for computer verification of EP for Lee weight on $R = \mathbb{Z}/N\mathbb{Z}$ for $N \leq 2048$.

Cases of maximal symmetry

- ▶ There has been progress on finding more explicit conditions over ring alphabets ($A = R$) when the weight w has maximal symmetry: $G_{lt} = G_{rt} = \mathcal{U}(R)$.
- ▶ When R is a product of chain rings: Greferath, Mc Fadden, Zumbrägel, 2013.
- ▶ When R is a principal ideal ring: Greferath, Honold, Mc Fadden, Wood, Zumbrägel, 2014. Here $\det \mathcal{A}$ is factored into terms $\sum_{0 < dR \leq aR} w(d) \mu(0, dR)$, for $a \in R$, where μ is the Möbius function for the poset of principal right ideals of R .

Examples over $\mathbb{Z}/N\mathbb{Z}$

- ▶ In addition to the Hamming weight, there are three additional weights that are easy to define on $\mathbb{Z}/N\mathbb{Z}$.
- ▶ Lee weight: viewing $\mathbb{Z}/N\mathbb{Z} = \{0, 1, \dots, N - 1\}$, Lee weight is $w_L(a) = \min\{a, N - a\}$.
- ▶ Euclidean weight: $w_E(a) = w_L(a)^2$.
- ▶ Complex Euclidean weight:
 $|\exp(2\pi ia/N) - 1|^2 = 2 - 2 \cos(2\pi a/N)$ (square of complex length).

Facts about EP over $\mathbb{Z}/N\mathbb{Z}$

- ▶ Only the complex Euclidean weight is easy: it is the egalitarian weight using $U = \{\pm 1\}$.
- ▶ EP for Lee weight has been numerically verified (\mathcal{A} invertible) for $N \leq 2048$. It has been proven for $N = 2^k, 3^k$ and for $N = p = 2q + 1$ or $p = 4q + 1$ (Barra) where p and q are prime.